

q -GENERATING FUNCTIONS FOR ONE AND TWO VARIABLES.

THOMAS ERNST

CONTENTS

1. Preliminaries	1
2. One variable	7
2.1. Special cases	10
3. Two variables	11
3.1. Special cases	15
References	18

ABSTRACT. We use a multidimensional extension of Bailey's transform to derive two very general q -generating functions, which are q -analogues of a paper by Exton [10]. These expressions are then specialised to give more practical formulae, which are q -analogues of generating relations for Karlssons generalised Kampé de Fériet function. A number of examples are given including q -Laguerre polynomials of two variables.

1. PRELIMINARIES

The purpose of this paper is to continue the study of q -special functions by the method outlined in [2]–[7]. The paper is a q -analogue of Exton [10]. All of Exton's results are obtained as the special case $q = 1$ by this method for $n \leq 2$.

We begin with a few definitions.

Definition 1. The power function is defined by $q^a = e^{a \log(q)}$. We always use the principal branch of the logarithm.

The q -analogues of a complex number a and of the factorial function are defined by:

$$(1) \quad \{a\}_q = \frac{1 - q^a}{1 - q}, \quad q \in \mathbb{C} \setminus \{1\},$$

$$(2) \quad \{n\}_q! = \prod_{k=1}^n \{k\}_q, \quad \{0\}_q! = 1, \quad q \in \mathbb{C},$$

Date: January 10, 2008.

⁰1991 Mathematics Subject Classification: Primary 33D70; Secondary 33C65

Definition 2. The q -hypergeometric series was developed by Heine 1846 as a generalization of the hypergeometric series:

$$(3) \quad {}_2\phi_1(a, b; c|q, z) = \sum_{n=0}^{\infty} \frac{\langle a; q \rangle_n \langle b; q \rangle_n}{\langle 1; q \rangle_n \langle c; q \rangle_n} z^n,$$

with the notation for the q -shifted factorial (compare [12, p.38])

$$(4) \quad \langle a; q \rangle_n = \begin{cases} 1, & n = 0; \\ \prod_{m=0}^{n-1} (1 - q^{a+m}) & n = 1, 2, \dots, \end{cases}$$

which is introduced in this paper.

Remark 1. The Watson notation [11] will also be used.

$$(5) \quad (a; q)_n = \begin{cases} 1, & n = 0; \\ \prod_{m=0}^{n-1} (1 - aq^m), & n = 1, 2, \dots, \end{cases}$$

Definition 3. Furthermore,

$$(6) \quad (a; q)_{\infty} = \prod_{m=0}^{\infty} (1 - aq^m), \quad 0 < |q| < 1.$$

$$(7) \quad (a; q)_{\alpha} = \frac{(a; q)_{\infty}}{(aq^{\alpha}; q)_{\infty}}, \quad a \neq q^{-m-\alpha}, m = 0, 1, \dots$$

Definition 4. In the following, $\frac{\mathbb{C}}{\mathbb{Z}}$ will denote the space of complex numbers mod $\frac{2\pi i}{\log q}$. This is isomorphic to the cylinder $\mathbb{R} \times e^{2\pi i\theta}$, $\theta \in \mathbb{R}$. The operator

$$\sim: \frac{\mathbb{C}}{\mathbb{Z}} \mapsto \frac{\mathbb{C}}{\mathbb{Z}}$$

is defined by

$$(8) \quad a \mapsto a + \frac{\pi i}{\log q}.$$

Furthermore we define

$$(9) \quad \widetilde{\langle a; q \rangle}_n = \langle \tilde{a}; q \rangle_n.$$

By (8) it follows that

$$(10) \quad \widetilde{\langle a; q \rangle}_n = \prod_{m=0}^{n-1} (1 + q^{a+m}),$$

where this time the tilde denotes an involution which changes a minus sign to a plus sign in all the n factors of $\langle a; q \rangle_n$.

The following simple rules follow from (8). Clearly the first two equations are applicable to q -exponents. Compare [24, p. 110].

$$(11) \quad \tilde{a} \pm b \equiv \widetilde{a \pm b} \pmod{\frac{2\pi i}{\log q}},$$

$$(12) \quad \tilde{a} \pm \tilde{b} \equiv a \pm b \pmod{\frac{2\pi i}{\log q}},$$

$$(13) \quad q^{\tilde{a}} = -q^a,$$

where the second equation is a consequence of the fact that we work mod $\frac{2\pi i}{\log q}$.

We will use the following abbreviation

$$(14) \quad \langle (a); q \rangle_n \equiv \langle a_1, \dots, a_A; q \rangle_n = \prod_{j=1}^A \langle a_j; q \rangle_n.$$

Definition 5. Generalizing Heine's series, we shall define a q -hypergeometric series by (compare [11, p.4]):

$$(15) \quad \begin{aligned} & {}_p\phi_r(\hat{a}_1, \dots, \hat{a}_p; \hat{b}_1, \dots, \hat{b}_r | q, z) \equiv {}_p\phi_r \left[\begin{matrix} \hat{a}_1, \dots, \hat{a}_p \\ \hat{b}_1, \dots, \hat{b}_r \end{matrix} \middle| q, z \right] = \\ & = \sum_{n=0}^{\infty} \frac{\langle \hat{a}_1, \dots, \hat{a}_p; q \rangle_n}{\langle 1, \hat{b}_1, \dots, \hat{b}_r; q \rangle_n} \left[(-1)^n q^{\binom{n}{2}} \right]^{1+r-p} z^n, \end{aligned}$$

where $q \neq 0$ when $p > r + 1$, and

$$(16) \quad \hat{a} = \begin{cases} a \\ \tilde{a} \end{cases}$$

We will skip the \hat{a} for the rest of the paper.

Definition 6. The following generalization of (15) will sometimes be used:

$$(17) \quad \begin{aligned} & {}_{p+p'}\phi_{r+r'}(a_1, \dots, a_p; b_1, \dots, b_r | q, z || s_1, \dots, s_{p'}; t_1, \dots, t_{r'}) \equiv \\ & {}_{p+p'}\phi_{r+r'} \left[\begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_r \end{matrix} \middle| q, z || \begin{matrix} s_1, \dots, s_{p'} \\ t_1, \dots, t_{r'} \end{matrix} \right] = \\ & = \sum_{n=0}^{\infty} \frac{\langle a_1; q \rangle_n \dots \langle a_p; q \rangle_n}{\langle 1; q \rangle_n \langle b_1; q \rangle_n \dots \langle b_r; q \rangle_n} \left[(-1)^n q^{\binom{n}{2}} \right]^{1+r+r'-p-p'} \times \\ & z^n \prod_{k=1}^{p'} (s_k; q)_n \prod_{k=1}^{r'} (t_k; q)_n^{-1}, \end{aligned}$$

where $q \neq 0$ when $p + p' > r + r' + 1$.

Remark 2. Equation (17) is used in certain special cases when we need factors $(t; q)_n$ in the q -series.

Definition 7. Let the q -Pochhammer symbol $\{a\}_{n,q}$ be defined by

$$(18) \quad \{a\}_{n,q} = \prod_{m=0}^{n-1} \{a+m\}_q.$$

An equivalent symbol is defined in [9, p.18] and is used throughout that book.

This quantity can be very useful in some cases where we are looking for q -analogues and it is included in the new notation.

If $|q| > 1$, or

$0 < |q| < 1$ and $|z| < |1 - q|^{-1}$, the q -exponential function $E_q(z)$ was defined by Jackson [13] 1904, and by Exton [9]

$$(19) \quad E_q(z) = \sum_{k=0}^{\infty} \frac{1}{\{k\}_q!} z^k.$$

Two q -analogues of the trigonometric functions are defined by

$$(20) \quad \text{Sin}_q(x) = \frac{1}{2i}(E_q(ix) - E_q(-ix)),$$

and

$$(21) \quad \text{Cos}_q(x) = \frac{1}{2}(E_q(ix) + E_q(-ix)).$$

The following multidimensional generalization of Bailey's transform was given by Exton [8, p.139].

Theorem 1.1. *If*

$$(22) \quad \gamma_{m_1, \dots, m_n} = \sum_{p_1=m_1, \dots, p_n=m_n}^{\infty} \delta_{p_1, \dots, p_n} u_{p_1-m_1, \dots, p_n-m_n} v_{p_1+m_1, \dots, p_n+m_n},$$

$$(23) \quad \beta_{m_1, \dots, m_n} = \sum_{p_1, \dots, p_n=0}^{m_1, \dots, m_n} \alpha_{p_1, \dots, p_n} u_{m_1-p_1, \dots, m_n-p_n} v_{p_1+m_1, \dots, p_n+m_n},$$

then formally

$$(24) \quad \sum_{\vec{m}} \alpha_{m_1, \dots, m_n} \gamma_{m_1, \dots, m_n} = \sum_{\vec{m}} \beta_{m_1, \dots, m_n} \delta_{m_1, \dots, m_n}.$$

We assume that α, δ, u, v are functions of m_1, \dots, m_n only. The notation $\sum_{\vec{m}}$ denotes a multiple summation with the indices m_1, \dots, m_n running over all non-negative integer values.

Definition 8. The following notation will be convenient.

$$(25) \quad \text{QE}(x) = q^x.$$

When there are several q :s, we generalize this to

$$(26) \quad \text{QE}(x, q_i) = q_i^x.$$

If $\{x_j\}_{j=1}^n$ and $\{y_j\}_{j=1}^n$ are two arbitrary sequences of complex numbers, then their scalar product is defined by

$$(27) \quad x\vec{y} = \sum_{j=1}^n x_j y_j.$$

We will only need one q -Lauricella function, which is defined by

$$(28) \quad \begin{aligned} & \Phi_D^{(n)}(a, b_1, \dots, b_n; c|q; x_1, \dots, x_n) = \\ & = \sum_{\vec{m}} \frac{\langle a; q \rangle_{m_1+\dots+m_n} \langle b_1; q \rangle_{m_1} \dots \langle b_n; q \rangle_{m_n} \prod_{j=1}^n x_j^{m_j}}{\langle c; q \rangle_{m_1+\dots+m_n} \prod_{j=1}^n \langle 1; q \rangle_{m_j}}. \end{aligned}$$

The following reduction theorem is a q -analogue of Appell and Kampé de Fériet [1, p. 116]. The proof uses one of the q -Vandermonde summation formulas. Because there are two such formulas, there is a quite similar equation, which was published for $n = 2$ in [22, p. 224, 4.18].

Theorem 1.2.

$$(29) \quad \begin{aligned} & \Phi_D^{(n)}(a, b_1, \dots, b_n; c|q; x, xq^{-b_2}, xq^{-b_2-b_3}, \dots, xq^{-b_2-\dots-b_n}) = \\ & = {}_2\phi_1(a, b_1 + \dots + b_n; c|q, xq^{-b_2-\dots-b_n}). \end{aligned}$$

Proof. In the LHS of (29) we change summation indices to $\{k_l\}_{l=1}^n$, where

$$(30) \quad k_l = \sum_{s=l}^n m_s.$$

By matrix inversion, this is equivalent to

$$(31) \quad m_l = k_l - k_{l+1}, \quad 1 \leq l \leq n-1, \quad m_n = k_n.$$

$$\begin{aligned}
(32) \quad LHS &= \sum_{m_i} \frac{\langle a; q \rangle_{m_1+\dots+m_n} \langle b_1; q \rangle_{m_1} \dots \langle b_n; q \rangle_{m_n} x^{m_1+\dots+m_n} \prod_{j=2}^n q^{-m_j(b_2+\dots+b_j)}}{\langle c; q \rangle_{m_1+\dots+m_n} \prod_{j=1}^n \langle 1; q \rangle_{m_j}} = \\
&= \sum_{k_i} \frac{\langle a; q \rangle_{k_1} \prod_{j=1}^{n-1} \langle b_j; q \rangle_{k_j-k_{j+1}} \langle b_n; q \rangle_{k_n} x^{k_1} \prod_{j=2}^{n-1} q^{(k_{j+1}-k_j)(b_2+\dots+b_j)} q^{(-k_n)(b_2+\dots+b_n)}}{\langle c; q \rangle_{k_1} \prod_{j=1}^{n-1} \langle 1; q \rangle_{k_j-k_{j+1}} \langle 1; q \rangle_{k_n}} \\
&= \sum_{k_i} \frac{\langle a; q \rangle_{k_1} \prod_{j=1}^{n-1} \langle b_j; q \rangle_{k_j} \langle -k_j; q \rangle_{k_{j+1}} q^{(k_{j+1})(-b_j+1)} \langle b_n; q \rangle_{k_n} x^{k_1}}{\langle c; q \rangle_{k_1} \prod_{j=1}^{n-1} \langle 1-b_j-k_j; q \rangle_{k_{j+1}} \langle 1; q \rangle_{k_j} \langle 1; q \rangle_{k_n}} \times \\
&\quad \prod_{j=2}^{n-1} q^{(k_{j+1}-k_j)(b_2+\dots+b_j)} q^{(-k_n)(b_2+\dots+b_n)} \\
&= \sum_{k_1, \dots, k_{n-1}} \frac{\langle a, b_1; q \rangle_{k_1} x^{k_1} \prod_{j=2}^{n-2} \langle b_j, -k_{j-1}; q \rangle_{k_j} q^{(k_j)(1-b_j-b_{j-1})}}{\langle c, 1; q \rangle_{k_1} \prod_{j=2}^{n-2} \langle 1, 1-b_{j-1}-k_{j-1}; q \rangle_{k_j}} \times \\
&\quad \frac{\langle b_{n-1}, -k_{n-2}, b_n+b_{n-1}; q \rangle_{k_{n-1}} q^{(k_{n-1})(1-b_{n-2}-b_{n-1}-b_n)}}{\langle 1, 1-b_{n-2}-k_{n-2}, b_{n-1}; q \rangle_{k_{n-1}}} = \\
&= \sum_{k_1, \dots, k_{n-2}} \frac{\langle a, b_1; q \rangle_{k_1} x^{k_1} \prod_{j=2}^{n-2} \langle b_j, -k_{j-1}; q \rangle_{k_j} q^{(k_j)(1-b_j-b_{j-1})}}{\langle c, 1; q \rangle_{k_1} \prod_{j=2}^{n-2} \langle 1, 1-b_{j-1}-k_{j-1}; q \rangle_{k_j}} \times \\
&\quad \frac{\langle 1-b_n-b_{n-1}-b_{n-2}-k_{n-2}; q \rangle_{k_{n-2}}}{\langle 1-b_{n-2}-k_{n-2}; q \rangle_{k_{n-2}}} = \\
&= \sum_{k_1, \dots, k_{n-2}} \frac{\langle a, b_1; q \rangle_{k_1} x^{k_1} \prod_{j=2}^{n-2} \langle b_j, -k_{j-1}; q \rangle_{k_j} q^{(k_j)(1-b_j-b_{j-1})}}{\langle c, 1; q \rangle_{k_1} \prod_{j=2}^{n-2} \langle 1, 1-b_{j-1}-k_{j-1}; q \rangle_{k_j}} \times \\
&\quad \frac{\langle b_n+b_{n-1}+b_{n-2}; q \rangle_{k_{n-2}} q^{-k_{n-2}(b_n+b_{n-1})}}{\langle b_{n-2}; q \rangle_{k_{n-2}}}.
\end{aligned}$$

We can continue this process to obtain the final result. \square

The following generalization of (29) is a q -analogue of [17] and [18]. Compare [20, (4.3), p. 107]

Theorem 1.3. *If $\{C_n\}_{n=0}^\infty$, $\{\alpha_n\}_{n=0}^\infty$ are sequences of arbitrary complex numbers then*

$$\begin{aligned}
(33) \quad &\sum_{\mathbf{m}} \frac{C_{m_1+\dots+m_n} \prod_{j=1}^n x^{m_j} \langle \alpha_j; q \rangle_{m_j}}{\prod_{j=1}^n \langle 1; q \rangle_{m_j}} \text{QE} \left(-\sum_{k=1}^n m_k \sum_{l=2}^k \alpha_l \right) = \\
&\sum_{N=0}^\infty \frac{C_N x^N \langle \sum_{k=1}^n \alpha_k; q \rangle_N}{\langle 1; q \rangle_N} \text{QE} \left(-N \sum_{l=2}^n \alpha_l \right).
\end{aligned}$$

Proof. We use induction. Suppose that (33) is true for $n > 1$, and denote the LHS by Δ_n , then

$$\begin{aligned}
 \Delta_{n+1} &= \sum_{m_{n+1}=0}^{\infty} \langle \alpha_{n+1}; q \rangle_{m_{n+1}} \frac{x^{m_{n+1}}}{\langle 1; q \rangle_{m_{n+1}}} \text{QE}(-m_{n+1}(\alpha_2 + \dots + \alpha_{n+1})) \times \\
 &\quad \sum_{m_1, \dots, m_n} C_{m_1 + \dots + m_{n+1}} \frac{x^{m_1 + \dots + m_n}}{\prod_{i=1}^n \langle 1; q \rangle_{m_i}} \prod_{k=1}^n \langle \alpha_k; q \rangle_{m_k} \text{QE} \left(-m_k \sum_{l=2}^k \alpha_l \right) \\
 (34) \quad &= \sum_{m_{n+1}=0}^{\infty} \langle \alpha_{n+1}; q \rangle_{m_{n+1}} \frac{x^{m_{n+1}}}{\langle 1; q \rangle_{m_{n+1}}} \text{QE} \left(-m_{n+1} \sum_{l=2}^{n+1} \alpha_l \right) \times \\
 &\quad \sum_{N=0}^{\infty} C_{N+m_{n+1}} \left\langle \sum_{k=1}^n \alpha_k; q \right\rangle_N \frac{x^N}{\langle 1; q \rangle_N} \text{QE} \left(-N \sum_{l=2}^n \alpha_l \right) \\
 &= \sum_{N=0}^{\infty} C_N x^N \text{QE} \left(-N \sum_{l=2}^{n+1} \alpha_l \right) \frac{\langle \sum_{k=1}^{n+1} \alpha_k; q \rangle_N}{\langle 1; q \rangle_N},
 \end{aligned}$$

where in the last step we used the induction hypothesis for $n = 2$, with the following values of the parameters in (33).

$$(35) \quad m_1 \rightarrow N, \quad m_2 \rightarrow m_n + 1, \quad \alpha_2 \rightarrow \alpha_n + 1, \quad x \rightarrow x \text{QE} \left(- \sum_{l=2}^n \alpha_l \right).$$

□

Remark 3. There is a dual too.

2. ONE VARIABLE

We begin with the case $n = 1$. In the rest of this paper, we assume that $|t| < 1$, $|q| < 1$.

Theorem 2.1. *If $C(m)$ is any arbitrary function, then, formally*

$$\begin{aligned}
 (36) \quad &\sum_m \frac{C(m) \langle d; q \rangle_m t^m (tq^{d+m}; q)_{\infty}}{(t; q)_{\infty}} = \\
 &= \sum_m \frac{\langle d; q \rangle_m t^m}{\langle 1; q \rangle_m} \sum_{p=0}^{\infty} C(p) \langle -m; q \rangle_p (-1)^p q^{mp - \binom{p}{2}}.
 \end{aligned}$$

Proof. In theorem 1.1 put

$$(37) \quad \alpha_m = C(m),$$

$$(38) \quad u_m = \frac{1}{\langle 1; q \rangle_m},$$

$$(39) \quad v_m = 1$$

and

$$(40) \quad \delta_m = \langle d; q \rangle_m t^m.$$

Now (22) and (23) imply that

$$(41) \quad \begin{aligned} \beta_m &= \sum_{p=0}^m \frac{C(p)}{\langle 1; q \rangle_{m-p}} = \\ &= \sum_{p=0}^m \frac{C(p) \langle -m; q \rangle_p}{\langle 1; q \rangle_m} (-1)^p q^{mp - \binom{p}{2}}. \end{aligned}$$

and

$$(42) \quad \begin{aligned} \gamma_m &= \sum_{p=m}^{\infty} \frac{\langle d; q \rangle_p t^p}{\langle 1; q \rangle_{p-m}} = \sum_{p=0}^{\infty} \frac{\langle d; q \rangle_{p+m} t^{p+m}}{\langle 1; q \rangle_p} = \\ &= \langle d; q \rangle_m t^m \sum_{p=0}^{\infty} \frac{\langle d+m; q \rangle_p t^p}{\langle 1; q \rangle_p} = \\ &= \langle d; q \rangle_m t^m {}_1\phi_0(d+m; -|q, t) \\ &= \langle d; q \rangle_m t^m \frac{(tq^{d+m}; q)_{\infty}}{(t; q)_{\infty}}. \end{aligned}$$

The proof is completed by substituting (41) and (42) into (24). \square

Theorem 2.2. *If $C(m)$ is any arbitrary function of m , then, formally*

$$(43) \quad \begin{aligned} E_q(t) \sum_m C(m) t^m (1-q)^m = \\ \sum_m \frac{t^m (1-q)^m}{\langle 1; q \rangle_m} \sum_{p=0}^{\infty} C(p) \langle -m; q \rangle_p (-1)^p q^{mp - \binom{p}{2}}. \end{aligned}$$

Proof. Let $d \rightarrow \infty$ in (36). \square

The theorems 2.1 and 2.2 are much too general for many practical purposes when deriving generating functions for various classes of q -hypergeometric polynomials. A more convenient form is obtained by considering the following special case.

$$(44) \quad C(m) = \frac{\langle (a), (f); q \rangle_m (-x)^m q^{\theta(m)}}{\langle (h), (g), 1; q \rangle_m},$$

where $\theta(m)$ is an arbitrary function.

Theorem 2.1 can be written as

$$\begin{aligned}
 & \sum_m \frac{\langle (a), (f), d; q \rangle_m (-x)^m t^m q^{\theta(m)}}{\langle (h), (g), 1; q \rangle_m (t; q)_{d+m}} = \\
 (45) \quad & = \sum_m \frac{\langle d; q \rangle_m t^m}{\langle 1; q \rangle_m} \sum_{p=0}^{\infty} \frac{\langle (a), (f); q \rangle_p (x)^p q^{\theta(p)}}{\langle (h), (g), 1; q \rangle_p} \times \\
 & \langle -m; q \rangle_p q^{mp - \binom{p}{2}}.
 \end{aligned}$$

Remark 4. (45) is a q -analogue of [21, p. 328, (1.2)] and closely resembles [23, p. 108, (3.33)].

The confluent form

$$\begin{aligned}
 & E_q(t) \sum_m \frac{\langle (a), (f); q \rangle_m (-x)^m}{\langle (h), (g), 1; q \rangle_m} t^m (1-q)^m q^{\theta(m)} = \\
 (46) \quad & = \sum_m \frac{t^m (1-q)^m}{\langle 1; q \rangle_m} \sum_{p=0}^{\infty} \frac{\langle (a), (f); q \rangle_p (x)^p q^{\theta(p)}}{\langle (h), (g), 1; q \rangle_p} \times \\
 & \langle -m; q \rangle_p q^{mp - \binom{p}{2}}
 \end{aligned}$$

follows similarly from theorem 2.2.

Another special case is obtained by putting

$$(47) \quad C(m) = \frac{\langle (a), (f), x; q \rangle_m (-1)^m q^{\theta(m)}}{\langle (h), (g), 1; q \rangle_m},$$

where $\theta(m)$ is an arbitrary function.

Theorem 2.1 can then be written as

$$\begin{aligned}
 & \sum_m \frac{\langle (a), (f), d, x; q \rangle_m (-1)^m t^m q^{\theta(m)}}{\langle (h), (g), 1; q \rangle_m (t; q)_{d+m}} = \\
 (48) \quad & = \sum_m \frac{\langle d; q \rangle_m t^m}{\langle 1; q \rangle_m} \sum_{p=0}^{\infty} \frac{\langle (a), (f), x; q \rangle_p q^{\theta(p)}}{\langle (h), (g), 1; q \rangle_p} \times \\
 & \langle -m; q \rangle_p q^{mp - \binom{p}{2}}.
 \end{aligned}$$

The confluent form

$$\begin{aligned}
 & E_q(t) \sum_m \frac{\langle (a), (f), x; q \rangle_m (-1)^m}{\langle (h), (g), 1; q \rangle_m} t^m (1-q)^m q^{\theta(m)} = \\
 (49) \quad & = \sum_m \frac{t^m (1-q)^m}{\langle 1; q \rangle_m} \sum_{p=0}^{\infty} \frac{\langle (a), (f), x; q \rangle_p q^{\theta(p)}}{\langle (h), (g), 1; q \rangle_p} \times \\
 & \langle -m; q \rangle_p q^{mp - \binom{p}{2}}
 \end{aligned}$$

follows similarly from theorem 2.2.

2.1. **Special cases.** Put $A = F = G = 0$, $H = 1$, $\theta(m) = m^2$. in (45) to obtain

$$\begin{aligned}
 (50) \quad & \sum_{m=0}^{\infty} \frac{(-x)^m}{\langle h; q \rangle_m \langle 1; q \rangle_m} \frac{\langle d; q \rangle_m t^m q^{m^2}}{(t; q)_{d+m}} = \\
 & = \sum_{m=0}^{\infty} \frac{\langle d; q \rangle_m t^m}{\langle 1; q \rangle_m} \sum_{p=0}^{\infty} \frac{x^p}{\langle h; q \rangle_p \langle 1; q \rangle_p} \times \\
 & \quad \langle -m; q \rangle_p q^{mp+p^2-\binom{p}{2}}.
 \end{aligned}$$

By a change of variables $x \rightarrow xq^{h-1}(1-q)$ this is equivalent to

$$\begin{aligned}
 (51) \quad & \sum_{n=0}^{\infty} \frac{\{c\}_{n,q} L_{n,q}^{(\alpha)}(x) t^n}{\{1+\alpha\}_{n,q}} = \sum_{n=0}^{\infty} \frac{\{c\}_{n,q} q^{n^2+\alpha n} (-xt)^n}{\{n\}_q! \{1+\alpha\}_{n,q} (t; q)_{c+n}} \\
 & \equiv \frac{1}{(t; q)_c} {}_1\phi_2(c; 1+\alpha|q; -xtq^{1+\alpha}(1-q)||-; tq^c).
 \end{aligned}$$

This is a wellknown generating function for the q -Laguerre polynomials.

Put $A = H = 1$, $F = G = 0$, $\theta(m) = \binom{m}{2}$ in (46). The little q -Jacobi polynomials are defined by

$$(52) \quad P_n(x; a, b|q) = \sum_{p=0}^n \frac{\langle a+b+n+1, -n; q \rangle_p x^p q^p}{\langle a+1, 1; q \rangle_p}.$$

Then we obtain the following generating function for the little q -Jacobi polynomials:

$$\begin{aligned}
 (53) \quad & \sum_n \frac{t^n (1-q)^n}{\langle 1; q \rangle_n} P_n(xq^{n-1}; a, b|q) \\
 & = E_q(t) {}_1\phi_1(a+b+n+1; a+1|q, xt(1-q)).
 \end{aligned}$$

Denote

$$(54) \quad {}_4\phi_7(\alpha) \equiv {}_4\phi_7 \left[\begin{matrix} \frac{a+b+n+1}{2}, \widetilde{\frac{a+b+n+1}{2}}, \frac{a+b+n+2}{2}, \widetilde{\frac{a+b+n+2}{2}} \\ \frac{1+a}{2}, \widetilde{\frac{1+a}{2}}, \frac{2+a}{2}, \widetilde{\frac{2+a}{2}}, \frac{1}{2}, \widetilde{\frac{1}{2}}, \widetilde{1} \end{matrix} \middle| q, q(1-q)^2 x^2 t^2 \right],$$

$$(55) \quad {}_4\phi_7(\beta) \equiv {}_4\phi_7 \left[\begin{matrix} \frac{a+b+n+2}{2}, \widetilde{\frac{a+b+n+2}{2}}, \frac{a+b+n+3}{2}, \widetilde{\frac{a+b+n+3}{2}} \\ \frac{2+a}{2}, \widetilde{\frac{2+a}{2}}, \frac{3+a}{2}, \widetilde{\frac{3+a}{2}}, \frac{3}{2}, \widetilde{\frac{3}{2}}, \widetilde{1} \end{matrix} \middle| q, q^3(1-q)^2 x^2 t^2 \right].$$

Making use of the decomposition of a series into even and odd parts from [19, p.200,208], we can rewrite (53) in the form

$$(56) \quad \sum_{n=0}^{\infty} \frac{P_{2n}(xq^{2n-1}; a, b|q)t^{2n}}{\{2n\}_q!} + \sum_{n=0}^{\infty} \frac{P_{2n+1}(xq^{2n}; a, b|q)t^{2n+1}}{\{2n+1\}_q!} \\ = E_q(t) \left[{}_4\phi_7(\alpha) - xt \frac{\{1+a+b+n\}_q}{\{1+a\}_q} {}_4\phi_7(\beta) \right],$$

and replacing t in (56) by it , we obtain

$$(57) \quad \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n} P_{2n}(xq^{2n-1}; a, b|q)}{\{2n\}_q!} + i \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n+1} P_{2n+1}(xq^{2n}; a, b|q)}{\{2n+1\}_q!} \\ = (Cos_q(t) + iSin_q(t)) \left[{}_4\phi_7(\alpha) - ixt \frac{\{1+a+b+n\}_q}{\{1+a\}_q} {}_4\phi_7(\beta) \right].$$

Next equate real and imaginary parts from both sides to arrive at the generating functions

$$(58) \quad \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n} P_{2n}(xq^{2n-1}; a, b|q)}{\{2n\}_q!} \\ = Cos_q(t) {}_4\phi_7(\alpha) + xt Sin_q(t) \frac{\{1+a+b+n\}_q}{\{1+a\}_q} {}_4\phi_7(\beta)$$

and

$$(59) \quad \sum_{n=0}^{\infty} \frac{(-1)^n t^{2n+1} P_{2n+1}(xq^{2n}; a, b|q)}{\{2n+1\}_q!} \\ = Sin_q(t) {}_4\phi_7(\alpha) - xt Cos_q(t) \frac{\{1+a+b+n\}_q}{\{1+a\}_q} {}_4\phi_7(\beta).$$

Finally put $A = F = G = 0$, $H = 1$, $\theta(m) = \binom{m}{2}$ in (48) to obtain the following generating function for q -Meixner polynomials from [23, p. 103, (3.10)]

$$(60) \quad \sum_m \frac{\langle x, d; q \rangle_m (-t)^m q^{\binom{m}{2}}}{\langle 1, h; q \rangle_m (t; q)_{d+m}} = \sum_m \frac{\langle d; q \rangle_m t^m}{\langle 1; q \rangle_m} {}_2\phi_1(x, -m_1; h|q, q^{m_1}).$$

3. TWO VARIABLES

We can generalize (36) to two variables.

Theorem 3.1. *If $C(m_1, m_2)$ is any arbitrary function of m_1, m_2 , then, formally*

$$\begin{aligned}
& \sum_{\vec{m}} \frac{C(m_1, m_2) \langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2}}{\langle k_1 + k_2; q \rangle_{m_1+m_2}} \times \\
& \frac{(tq^{d+m_1-k_2}; q)_\infty}{(tq^{-k_2-m_2}; q)_\infty} = \\
(61) \quad & = \sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} q^{\frac{m_2^2}{2}}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \times \\
& \sum_{p_1, p_2=0}^{\infty} C(p_1, p_2) \prod_{j=1}^2 \langle -m_j; q \rangle_{p_j} (-1)^{p_1+p_2} \text{QE} \left(- \sum_{j=1}^2 \binom{p_j}{2} \right. \\
& \left. - k_2(m_2 - p_2) + m_1 p_1 + p_2^2 \right).
\end{aligned}$$

Proof. In theorem 1.1 put

$$(62) \quad \alpha_{m_1, m_2} = C(m_1, m_2),$$

$$(63) \quad u_{m_1, m_2} = \frac{\text{QE}(\frac{3}{4}m_2^2 - k_2m_2)}{\prod_{j=1}^2 \langle 1; q \rangle_{m_j}},$$

$$(64) \quad v_{m_1, m_2} = q^{\frac{1}{4}m_2^2}$$

and

$$(65) \quad \delta_{m_1, m_2} = \frac{q^{-m_2^2} \langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2}}{\langle k_1 + k_2; q \rangle_{m_1+m_2}}.$$

Now (22) and (23) imply that

$$\begin{aligned}
(66) \quad \beta_{m_1, m_2} &= \sum_{p_1, p_2=0}^{m_1, m_2} \frac{C(p_1, p_2)}{\prod_{j=1}^2 \langle 1; q \rangle_{m_j - p_j}} \\
&\times \text{QE}(\frac{3}{4}(m_2 - p_2)^2 - k_2(m_2 - p_2) + \frac{1}{4}(m_2 + p_2)^2) = \\
&= \sum_{p_1, p_2=0}^{m_1, m_2} \frac{C(p_1, p_2) \prod_{j=1}^2 \langle -m_j; q \rangle_{p_j}}{\prod_{j=1}^2 \langle 1; q \rangle_{m_j}} (-1)^{p_1+p_2} \text{QE} \left(- \sum_{j=1}^2 \binom{p_j}{2} \right) \\
&\text{QE}(-k_2(m_2 - p_2) + m_2^2 + m_1 p_1 + p_2^2),
\end{aligned}$$

and

$$\begin{aligned}
 \gamma_{m_1, m_2} &= \sum_{p_1=m_1, p_2=m_2}^{\infty} \frac{\langle d; q \rangle_{p_1+p_2} \prod_{j=1}^2 \langle k_j; q \rangle_{p_j} t^{p_1+p_2}}{\prod_{j=1}^2 \langle 1; q \rangle_{p_j-m_j} \langle k_1+k_2; q \rangle_{p_1+p_2}} \\
 &\times \text{QE}(-p_2^2 + \frac{3}{4}(p_2 - m_2)^2 - k_2(p_2 - m_2) + \frac{1}{4}(m_2 + p_2)^2) = \\
 &= \sum_{p_1, p_2=0}^{\infty} \frac{\langle d; q \rangle_{p_1+m_1+p_2+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{p_j+m_j} t^{p_1+p_2+m_1+m_2}}{\prod_{j=1}^2 \langle 1; q \rangle_{p_j} \langle k_1+k_2; q \rangle_{p_1+p_2+m_1+m_2}} \\
 &\times \text{QE}(-p_2(k_2 + m_2)) = \\
 &= \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2}}{\langle k_1+k_2; q \rangle_{m_1+m_2}} \sum_{p_1, p_2=0}^{\infty} \text{QE}(-p_2(k_2 + m_2)) \times \\
 (67) \quad &\frac{\langle d + m_1 + m_2; q \rangle_{p_1+p_2} \prod_{j=1}^2 \langle k_j + m_j; q \rangle_{p_j} t^{p_1+p_2}}{\prod_{j=1}^2 \langle 1; q \rangle_{p_j} \langle k_1+k_2+m_1+m_2; q \rangle_{p_1+p_2}} = \\
 &= \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2}}{\langle k_1+k_2; q \rangle_{m_1+m_2}} \Phi_D^{(2)}(d + m_1 + m_2, \\
 &, k_1 + m_1, k_2 + m_2; k_1 + m_1 + k_2 + m_2 | q; t, tq^{-k_2-m_2}) \\
 &= \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2}}{\langle k_1+k_2; q \rangle_{m_1+m_2}} \times \\
 &{}_1\phi_0(d + m_1 + m_2; -|q, tq^{-k_2-m_2}) \\
 &= \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2}}{\langle k_1+k_2; q \rangle_{m_1+m_2}} \frac{1}{(tq^{-k_2-m_2}; q)_{d+m_1+m_2}}.
 \end{aligned}$$

The proof is completed by substituting (66) and (67) into (24). \square

Theorem 3.2. *If $C(m_1, m_2)$ is any arbitrary function of m_1, m_2 , then, formally*

$$\begin{aligned}
 (68) \quad &\sum_{\vec{m}} \frac{E_q(tq^{-k_2-m_2}) C(m_1, m_2) \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} (1-q)^{m_1+m_2}}{\langle k_1+k_2; q \rangle_{m_1+m_2}} = \\
 &= \sum_{\vec{m}} \frac{\prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} (1-q)^{m_1+m_2} q^{\frac{m_2^2}{2}}}{\langle k_1+k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \times \\
 &\sum_{p_1, p_2=0}^{\infty} C(p_1, p_2) \prod_{j=1}^2 \langle -m_j; q \rangle_{p_j} (-1)^{p_1+p_2} \text{QE} \left(-\sum_{j=1}^2 \binom{p_j}{2} \right) \\
 &\text{QE}(-k_2(m_2 - p_2) + m_1 p_1 + p_2^2).
 \end{aligned}$$

Proof. Let $d \rightarrow \infty$ in (61). \square

The theorems 3.1 and 3.2 are much too general for many practical purposes when deriving generating functions for various classes of hypergeometric polynomials. A more convenient form is obtained by considering the following special case.

$$(69) \quad C(m_1, m_2) = \frac{\langle (a); q \rangle_{m_1+m_2} q^{\theta(m_1, m_2)} \prod_{j=1}^2 \langle (f_j); q \rangle_{m_j} (-x_j)^{m_j}}{\langle (h); q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle (g_j), 1; q \rangle_{m_j}},$$

where $\theta(m_1, m_2)$ is an arbitrary function.

Theorem 3.1 can be written as

$$(70) \quad \begin{aligned} & \sum_{\vec{m}} \frac{\langle (a); q \rangle_{m_1+m_2} q^{\theta(m_1, m_2)} \prod_{j=1}^2 \langle (f_j); q \rangle_{m_j} (-x_j)^{m_j}}{\langle (h), k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle (g_j), 1; q \rangle_{m_j}} \times \\ & \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} (tq^{d+m_1-k_2}; q)_\infty}{(tq^{-k_2-m_2}; q)_\infty} = \\ & = \sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} q^{\frac{m_2^2}{2}}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \sum_{p_1, p_2=0}^{\infty} \\ & \frac{\langle (a); q \rangle_{p_1+p_2} q^{\theta(p_1, p_2)} \prod_{j=1}^2 \langle (f_j), -m_j; q \rangle_{p_j} x_j^{p_j}}{\langle (h); q \rangle_{p_1+p_2} \prod_{j=1}^2 \langle (g_j), 1; q \rangle_{p_j}} \times \\ & \text{QE} \left(-k_2(m_2 - p_2) + m_1 p_1 + p_2^2 - \sum_{j=1}^2 \binom{p_j}{2} \right). \end{aligned}$$

The following confluent form follows similarly from theorem (3.2).

$$(71) \quad \begin{aligned} & \sum_{\vec{m}} \frac{E_q(tq^{-k_2-m_2}) \langle (a); q \rangle_{m_1+m_2} q^{\theta(m_1, m_2)}}{\langle (h), k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle (g_j), 1; q \rangle_{m_j}} \times \\ & t^{m_1+m_2} (1-q)^{m_1+m_2} \prod_{j=1}^2 \langle (f_j), k_j; q \rangle_{m_j} (-x_j)^{m_j} = \\ & = \sum_{\vec{m}} \frac{\prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} (1-q)^{m_1+m_2} q^{\frac{m_2^2}{2}}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \sum_{p_1, p_2=0}^{\infty} \\ & \frac{\langle (a); q \rangle_{p_1+p_2} q^{\theta(p_1, p_2)} \prod_{j=1}^2 \langle (f_j), -m_j; q \rangle_{p_j} x_j^{p_j}}{\langle (h); q \rangle_{p_1+p_2} \prod_{j=1}^2 \langle (g_j), 1; q \rangle_{p_j}} \times \\ & \text{QE} \left(-k_2(m_2 - p_2) + m_1 p_1 + p_2^2 - \sum_{j=1}^2 \binom{p_j}{2} \right). \end{aligned}$$

3.1. **Special cases.**

Put $A = F = G = 0$, $H = 1$, $\theta(m_1, m_2) = m_1^2$ in (70). Then

$$(72) \quad \sum_{\vec{m}} \frac{q^{m_1^2} \langle d; q \rangle_{m_1+m_2} t^{m_1+m_2} \prod_{j=1}^2 (-x_j)^{m_j} \langle k_j; q \rangle_{m_j}}{\langle h, k_1 + k_2; q \rangle_{m_1+m_2} (tq^{-k_2-m_2}; q)_{d+m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} =$$

$$\sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} q^{\frac{m_2^2}{2}}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \sum_{p_1, p_2=0}^{m_1, m_2} \frac{\text{QE}(-k_2(m_2 - p_2))}{\langle h; q \rangle_{p_1+p_2}} \times$$

$$\frac{\prod_{j=1}^2 \langle -m_j; q \rangle_{p_j} (x_j)^{p_j}}{\prod_{j=1}^2 \langle 1; q \rangle_{p_j}} \text{QE} \left(m_1 p_1 + \sum_{j=1}^2 - \binom{p_j}{2} + p_j^2 \right).$$

By a change of variables $x_1 \rightarrow x_1 q^{h-1} (1 - q)$, $x_2 \rightarrow x_2 (1 - q)$ this is equivalent to (compare [10, A.17])

$$(73) \quad \sum_{\vec{m}} \frac{q^{m_1^2+m_1(h-1)} \langle d; q \rangle_{m_1+m_2} t^{m_1+m_2} \prod_{j=1}^2 (-x_j)^{m_j} \langle k_j; q \rangle_{m_j} (1 - q)^{m_1+m_2}}{\langle h, k_1 + k_2; q \rangle_{m_1+m_2} (tq^{-k_2-m_2}; q)_{d+m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}}$$

$$= \sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2}}{\langle k_1 + k_2, h; q \rangle_{m_1+m_2}} L_{m_1, m_2, k_2, q}^{h-1}(x_1, x_2),$$

where $L_{m_1, m_2, k_2, q}^\alpha(x_1, x_2)$ is the q -Laguerre polynomial in two variables given by

$$(74) \quad L_{m_1, m_2, k_2, q}^\alpha(x_1, x_2) = \frac{\langle \alpha + 1; q \rangle_{m_1+m_2}}{\prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \sum_{p_1, p_2=0}^{m_1, m_2} \frac{q^{p_1^2+\alpha p_1} \prod_{j=1}^2 \langle -m_j; q \rangle_{p_j} (x_j)^{p_j}}{\langle 1 + \alpha; q \rangle_{p_1+p_2} \prod_{j=1}^2 \langle 1; q \rangle_{p_j}}$$

$$\text{QE}(m_1 p_1 + \frac{1}{2} m_2^2 + p_2^2 - k_2(m_2 - p_2))(1 - q)^{p_1+p_2} \prod_{j=1}^2 \text{QE} \left(- \binom{p_j}{2} \right).$$

By letting $d = h$, $d = k_1 + k_2$ and $d \rightarrow \infty$ in (73), we obtain q -analogues of [10, A19-A21] for two variables.

Put $F = G = H = 0$, $A = 1$, $\theta(m_1, m_2) = m_1^2$ in (70). Then

$$\begin{aligned}
(75) \quad & \sum_{\vec{m}} \frac{q^{m_1^2} \langle a, d; q \rangle_{m_1+m_2} t^{m_1+m_2} \prod_{j=1}^2 (-x_j)^{m_j} \langle k_j; q \rangle_{m_j}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} (tq^{-k_2-m_2}; q)_{d+m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \cong \\
& \sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} q^{\frac{m_2^2}{2}}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \sum_{p_1, p_2=0}^{m_1, m_2} \text{QE}(-k_2(m_2 - p_2)) \times \\
& \frac{\langle a; q \rangle_{p_1+p_2} \prod_{j=1}^2 \langle -m_j; q \rangle_{p_j} (x_j)^{p_j}}{\prod_{j=1}^2 \langle 1; q \rangle_{p_j}} \text{QE} \left(m_1 p_1 + \sum_{j=1}^2 -\binom{p_j}{2} + p_j^2 \right) \cong \\
& \sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} t^{m_1+m_2} q^{\frac{m_2^2}{2}} \langle a; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} (-x_j)^{m_j}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \sum_{p_1, p_2=0}^{m_1, m_2} \\
& \frac{\prod_{j=1}^2 \langle -m_j; q \rangle_{p_j} (-x_j)^{-p_j}}{\langle -a + 1 - m_1 - m_2; q \rangle_{p_1+p_2} \prod_{j=1}^2 \langle 1; q \rangle_{p_j}} \text{QE}(-k_2(p_2) - m_1 p_1 + m_1^2 \\
& + p_1 p_2 - (p_1 + p_2)(m_1 + m_2) + \sum_{j=1}^2 p_j^2 - (a-1)p_j).
\end{aligned}$$

This is a q -analogue of the corrected form of [10, A22] for two variables. The symbol \cong denotes that the equality is purely formal. Put $A = H = 0$, $F = G = 1$, $\theta(m_1, m_2) = -m_2^2 + \sum_{j=1}^2 \binom{m_j}{2}$ in (70). Then (compare [10, A29])

$$\begin{aligned}
(76) \quad & \sum_{\vec{m}} \frac{q^{-m_2^2} \prod_{j=1}^2 \langle f_j; q \rangle_{m_j} (-x_j)^{m_j} q^{\binom{m_j}{2}}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle g_j, 1; q \rangle_{m_j}} \times \\
& \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} (tq^{d+m_1-k_2}; q)_\infty}{(tq^{-k_2-m_2}; q)_\infty} = \\
& = \sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} q^{\frac{m_2^2}{2} - k_2 m_2}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} {}_2\phi_1(f_1, -m_1; g_1 | q, x_1 q^{m_1}) \times \\
& {}_2\phi_1(f_2, -m_2; g_2 | q, x_2 q^{k_2}).
\end{aligned}$$

By letting $k_i = g_i$, $d = g_1 + g_2$, $d \rightarrow \infty$ and $(k_i = g_i, d \rightarrow \infty)$ in (76), we obtain q -analogues of [10, A30-A33].

Put $A = H = G = 0$, $F = 1$, $\theta(m_1, m_2) = -m_2^2 + \sum_{j=1}^2 \binom{m_j}{2}$ in (70).

Then (compare [10, A34])

$$\begin{aligned}
 & \sum_{\vec{m}} \frac{q^{-m_2^2} \prod_{j=1}^2 \langle f_j; q \rangle_{m_j} (-x_j)^{m_j} q^{\binom{m_j}{2}}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \times \\
 & \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} (tq^{d+m_1-k_2}; q)_\infty}{(tq^{-k_2-m_2}; q)_\infty} = \\
 (77) \quad & = \sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j} t^{m_1+m_2} q^{\frac{m_2^2}{2}-k_2m_2}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} {}_2\phi_1(f_1, -m_1; \infty | q, x_1 q^{m_1}) \times \\
 & {}_2\phi_1(f_2, -m_2; \infty | q, x_2 q^{k_2}).
 \end{aligned}$$

By letting $d \rightarrow \infty$ in (77), we obtain a q -analogue of [10, A36].

Put $A = F = H = 0$, $G = 1$, $\theta(m_1, m_2) = m_1^2$ in (70). Then (compare [10, A38])

$$\begin{aligned}
 & \sum_{\vec{m}} \frac{q^{m_1^2} \langle d; q \rangle_{m_1+m_2} t^{m_1+m_2} \prod_{j=1}^2 (-x_j)^{m_j} \langle k_j; q \rangle_{m_j}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} (tq^{-k_2-m_2}; q)_{d+m_1+m_2} \prod_{j=1}^2 \langle g_j, 1; q \rangle_{m_j}} = \\
 (78) \quad & \sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} t^{m_1+m_2} q^{\frac{m_2^2}{2}} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \sum_{p_1, p_2=0}^{m_1, m_2} \text{QE}(-k_2(m_2 - p_2)) \times \\
 & \frac{\prod_{j=1}^2 \langle -m_j; q \rangle_{p_j} (x_j)^{p_j}}{\prod_{j=1}^2 \langle g_j, 1; q \rangle_{p_j}} \text{QE}(m_1 p_1 + \sum_{j=1}^2 -\binom{p_j}{2} + p_j^2).
 \end{aligned}$$

By a change of variables $x_j \rightarrow x_j q^{g_j-1} (1-q)$, $j = 1, 2$ this is equivalent to

$$\begin{aligned}
 & \sum_{\vec{m}} \frac{q^{m_1^2} \langle d; q \rangle_{m_1+m_2} t^{m_1+m_2} (1-q)^{m_1+m_2} \prod_{j=1}^2 (-x_j)^{m_j} q^{m_j(g_j-1)} \langle k_j; q \rangle_{m_j}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} (tq^{-k_2-m_2}; q)_{d+m_1+m_2} \prod_{j=1}^2 \langle g_j, 1; q \rangle_{m_j}} \\
 & = \sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} t^{m_1+m_2} q^{\frac{m_2^2}{2}} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle 1; q \rangle_{m_j}} \sum_{p_1, p_2=0}^{m_1, m_2} \text{QE}(-k_2(m_2 - p_2)) \times \\
 (79) \quad & \frac{(1-q)^{p_1+p_2} \prod_{j=1}^2 \langle -m_j; q \rangle_{p_j} (x_j)^{p_j} q^{p_j(g_j-1)}}{\prod_{j=1}^2 \langle g_j, 1; q \rangle_{p_j}} \text{QE}(m_1 p_1 + \sum_{j=1}^2 -\binom{p_j}{2} + p_j^2) \\
 & = \sum_{\vec{m}} \frac{\langle d; q \rangle_{m_1+m_2} t^{m_1+m_2} q^{\frac{m_2^2}{2}-k_2m_2} \prod_{j=1}^2 \langle k_j; q \rangle_{m_j}}{\langle k_1 + k_2; q \rangle_{m_1+m_2} \prod_{j=1}^2 \langle g_j; q \rangle_{m_j}} L_{m_1, q}^{g_1-1}(x_1) L_{m_2, q}^{g_2-1}(x_2 q^{k_2-m_2}).
 \end{aligned}$$

By letting $k_i = g_i$, $d = g_1 + g_2$, $d \rightarrow \infty$ and $(k_i = g_i, d \rightarrow \infty)$ in (79), we obtain q -analogues of [10, A39-A42] for two variables.

Acknowledgments. I want to thank Per Karlsson who gave some valueable comments on the new method for q -hypergeometric series, and who told me about Exton's paper.

REFERENCES

- [1] Appell P and Kampé de Fériet, J, *Fonctions hypergéométriques et hypersphériques*, Paris 1926.
- [2] Ernst T, *The history of q -calculus and a new method*, U. U. D. M. Report 2000:16, ISSN 1101-3591, Department of Mathematics, Uppsala University, 2000.
- [3] Ernst T, *A new method and its Application to Generalized q -Bessel Polynomials*, U. U. D. M. Report 2001:10, ISSN 1101-3591, Department of Mathematics, Uppsala University, 2001.
- [4] Ernst T, *A new method for q -calculus* Uppsala dissertations in mathematics **25** 2002.
- [5] Ernst T, A method for q -calculus. *J. nonlinear Math. Physics* **10** No.4 (2003), 487-525.
- [6] Ernst T, Some results for q -functions of many variables. Preprint 2003.
- [7] Ernst T, q -Analogues of some operational formulas. Preprint 2003.
- [8] Exton H, Multiple hypergeometric functions and applications. Ellis Horwood, 1976.
- [9] Exton H, *q -Hypergeometric functions and applications*, Ellis Horwood, 1983.
- [10] Exton H, Two new multivariable generating relations, *Ark. Mat.* **30** (1992), no. 2, 245–258.
- [11] Gasper, G and Rahman M, *Basic hypergeometric series*, Cambridge, 1990.
- [12] Gelfand I M and Graev M I and Retakh V S, General hypergeometric systems of equations and series of hypergeometric type, *Russian Math. Surveys* **47** (1992), no. 4, 1–88.
- [13] Jackson F H, A basic-sine and cosine with symbolical solution of certain differential equations. *Proc. Edinburgh math. Soc.* **22** (1904), 28–39.
- [14] Jackson F H, On basic double hypergeometric functions, *Quart. J. Math., Oxford Ser.* **13** (1942), 69–82.
- [15] Jackson F H, Basic double hypergeometric functions, *Quart. J. Math., Oxford Ser.* **15**(1944), 49–61.
- [16] Hahn W, Beiträge zur Theorie der Heineschen Reihen, *Mathematische Nachrichten* **2** (1949), 340–379.
- [17] Panda R., Some multiple series transformations. *J nānabha Sect.* **A 4** (1974) 165–168.
- [18] Singhal B.M., On the reducibility of Lauricella's function F_D . *J nānabha* **A 4** (1974) 163–164.
- [19] Srivastava H M and Manocha H L, *A treatise on generating functions*, Ellis Horwood Series: John Wiley & Sons, Inc. , New York, 1984.
- [20] Srivastava H. M., A note on certain summation theorems for multiple hypergeometric series. *Simon Stevin* **52** (1978), no. 3 97–109.
- [21] Srivastava H. M., A family of q -generating functions. *Bull. Inst. Math. Acad. Sinica* 12 (1984), **4**, 327–336.
- [22] Srivastava H M and Jain V K, q -series identities and reducibility of basic double hypergeometric functions. *Canad. J. Math.* **38**, no. 1, (1986), 215–231.
- [23] Srivastava H. M. and Agarwal A. K., Generating functions for a class of q -polynomials. *Ann. Mat. Pura Appl.* (4) **154** (1989), 99–109.
- [24] Thomae J, Les séries Heineennes supérieures. *Ann. Mat. pura appl. Bologna*, II **4** (1871) 105–139.

DEPARTMENT OF MATHEMATICS, UPPSALA UNIVERSITY, P.O. BOX 480, SE-751 06 UPPSALA, SWEDEN

E-mail address: Thomas.Ernst@math.uu.se